



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

PETROGRAPHY.

Experimental Petrography. — Morozewicz¹ has just published a long paper on "Experimental investigations upon the formation of minerals in magmas" that will unquestionably take a place among the most important contributions to experimental geology that have been made within recent years. The author fused known mixtures of the various rock-producing compounds in a glass-furnace and thoroughly studied the resulting products. The conclusions reached by him are full of suggestiveness. The way seems to have been opened for a long line of important investigations to follow, some of which have already been entered upon.

The details of the experiments cannot be entered upon here, but some of the conclusions arrived at may be briefly indicated.

1. The structures of cooled magmas appear to depend upon exterior conditions of crystallization and upon their chemical composition, quantitative as well as qualitative.

2. The order of crystallization is determined by no one condition, such as fusibility, acidity, etc., but it depends upon a number of variable conditions, one of the most important of which is the quantity of the various constituents present as compared with their solubility in the molten mass. The ability of a substance to supersaturate the magma depends primarily upon its nature, the nature of the other substances composing the magma, and its temperature.

3. So far as the experiments touch upon the question of the differentiation of magmas, they seem to indicate that a molten mass may separate into layers or parts differing in density, and that this difference may be due to the fact that the bases FeO, MgO, CaO, separate as silicates by crystallization earlier than the remaining constituents.

Two pounds of granite, with the composition given below (I), were heated for five days. The mixture yielded a mass of glass which in its upper portion contained unmelted quartz grains and a considerable quantity of tridymite. Though the glass between the quartz grains in the upper portion of the crucible presented the same appearance as that in the lower parts, the composition of pieces taken from the two parts was found to be quite different. Under (II) we

¹ *Min. u. Petrog. Mitth.*, vol. xviii, pp. 1 and 105.

have the analysis of the glass from the upper part, and under (III) that of the glass from the lower portion.

	SiO ₂	Al ₂ O ₃	Fe ₂ O ₃	CaO	MgO	K ₂ O	Na ₂ O		Total.	Sp. Gr.
(I)	68.9	19.7	1.4	1.2	1.0	2.7	4.7	=	99.6	2.716
(II)	73.65	14.08	2.33	1.94	.65	2.61	3.86	=	99.12	2.2384
(III)	59.20	22.30	3.83	3.71	1.23	3.26	5.40	=	98.93	2.484

The Basic Rocks of Ivrea. — The basis rocks in the neighborhood of Ivrea, on the south side of the Alps, are shown by Schaefer¹ to be the result of cooling of a single magma. This yielded norites, diorites, gabbros, peridotites and both basic and acid dyke-rocks. The norites include hornblendic varieties, and the diorites, bronzitic, hornblendic, and biotitic phases. All these rocks have been subjected to the action of mountain-making forces. The norites have become schistose without suffering any essential mineralogical change. Some of the diorites have simply been made schistose, others have undergone a further change in that their dark, compact hornblende has passed over into a light green amphibole, while a final stage of alteration is represented by green schists, composed of zoisite, plagioclase, actinolite, chlorite, and epidote.

The dyke rocks cut the large basic masses and are always closely related to them chemically. The principal types are a labradorite (Labradorfels) and a fine-grained black rock which the author calls valbellite. This is made up of bronzite, olivine, and brown hornblende with pyrrhotite, spinel, and magnetite as accessories.

The Basalts of Steiermark. — Sigmund's² studies on the basalts of Steiermark are continued in an article in which are described the magma-basalts and basalt-tuffs of Fürstenfeld and the feldspar basalt of Weitendorf. The composition of the magma-basalt is shown by the figures below.

SiO ₂	TiO ₂	Fe ₂ O ₃	FeO	Al ₂ O ₃	CaO	MgO	Na ₂ O	K ₂ O	CO ₂	H ₂ O	Total.
46.76	tr.	5.33	5.62	17.93	8.24	7.31	3.53	2.20	1.33	1.83	= 100.08

Petrographical Notes. — Reinisch³ has found a specimen of teschenite in the museum at Minussinsk. It is labeled as having come from east of the salt lake Staniza on the river Bjelyi-Jjuss, Minussinsk parish, Jenisseisk gouvernement, East Siberia. It resembles very closely the West Carpathian rock. Among the other specimens from

¹ *Min. u. Petrog. Mitth.*, vol. xvii, p. 495.

² *Ibid.*, p. 256.

³ *Ibid.*, vol. xviii, p. 92.